

**Gebrauchsinweisung****Produktbeschreibung**

PermaFoam Classic ist ein Schaumverband, bestehend aus einem hydrophilen Polymer in dreidimensionaler Struktur, der wirksam Wundexsudat aufnimmt und so für ein feuchtes Wundmilieu sorgt. PermaFoam Classic Border verfügt zusätzlich über einen umlaufenden, selbsthaftenden Kleberand, der mit einem Polyacrylatkleber beschichtet ist.

**Zusammensetzung**

- PU-Schaum
- Polyacrylat
- PU-Folie

**Indikationen**

PermaFoam Classic ist ein steriler, wickelfreier Polyurethan-Schaumverband für den Einmalgebrauch, der bei moderat bis stark exsudierenden, akuten und chronischen Wunden indiziert ist, wie etwa venösen Ulzera (Beinulzera), Dekubitus Grad II–III und diabetischen Fußulzera. Gewebsnahtnahtstellen, Hautschürfwunden und Schnittwunden.

**Anwendungshinweise**

- Die Wunde mit physiologischer Lösung oder nach ärztlicher Anweisung reinigen und die umgebende Haut behutsam trocknen.
  - Die Wundfläche entsprechend der Wundgröße auswählen, sodass die mittige Kompressse die Wundränder etwa 1–2 cm überdeckt.
  - Für PermaFoam Classic Border: Die Schutzfolien abziehen, die weiße Seite von PermaFoam Classic auf die Wunde legen, sodass die beige Außenseite der Wunde abgewandt ist. Darüber einen zusätzlichen Fixierverband anlegen.
  - Für PermaFoam Classic Border: Die Schutzfolien abziehen, die weiße Seite der Schaumkompressse auf die Wunde legen und den Kleberand gut andrücken.
- PermaFoam Classic kann bis zu 7 Tage auf der Wunde verbleiben. Wie häufig ein Verbandwechsel erforderlich ist, hängt vom Wundzustand und der Exsudatmenge ab.

**PermaFoam Classic Sacral:** Siehe Diagramm 1.  
**PermaFoam Classic Concave:** Siehe Diagramm 2.  
**PermaFoam Classic Tracheostomy:** Siehe Diagramm 3.  
**PermaFoam Classic Border:** Siehe Diagramm 4.

**Kontraindikationen**

PermaFoam Classic sollte nicht angewendet werden, wenn eine Überempfindlichkeit gegenüber einem der Inhaltsstoffe vorliegt.

**Vorsichtsmaßnahmen**

- Das Produkt darf nicht bei infizierten Wunden verwendet werden.
- Bei Auftreten von Rötung oder allergischen Reaktionen die Behandlung mit PermaFoam Classic abbrechen.
- Bei Wunden infolge von Durchblutungsstörungen oder diabetischem Fußulkus sollte der Wundzustand täglich überprüft werden.
- Vor einer Behandlung mit Strahlung, einschließlich Röntgenstrahlung, Ultraschall und Mikrowellen wird vor einer Thermotherapie sollte das Produkt abgenommen werden.
- PermaFoam Classic darf nicht im Verbindung mit Wasserstoffperoxid verwendet werden.

**Produktentwurf**

Um das Risiko potentieller Infektionsgefahren oder einer Umweltverschmutzung zu minimieren, sollen Wegwerfkomponenten von PermaFoam Classic gemäß den geltenden lokalen Gesetzen, Vorschriften und Bestimmungen sowie den Standards zur Infektionsprävention entsorgt werden.

**Besondere Vorsichtsmaßnahmen**

- Es sind keine Daten verfügbar, die die Anwendung des Verbands bei sensiblen Patientengruppen wie Säuglingen, Kindern, Schwangeren oder stillenden Frauen unterstützen, und es liegen ebenfalls keine Daten vor, die dem widersprechen. Der Wundverband sollte bei diesen Patientengruppen daher nur mit Vorsicht und auf Empfehlung des Produkterzeugers verwendet werden.
- Hinweis für Anwender und/oder Patienten: Jedes schwerwiegende Vorkommnis im Zusammenhang mit dem Medizinprodukt muss dem Hersteller und der zuständigen Behörde des Mitgliedsstaats, in dem der Anwender und/oder Patient ansässig ist, gemeldet werden.

Wiederverwendung eines Einmalprodukts ist gefährlich. Die Wiederverwendung von Einmalprodukten, um sie erneut zu verwenden, kann ihre Integrität und Leistung deutlich beeinträchtigen.

\* Wenn PermaFoam Classic im folgenden Text genannt wird, bezieht sich dies auf alle Aufmachungen; davon abweichende Ausnahmen sind ausdrücklich erwähnt. Weitere Informationen auf Anfrage.

Stand of Information: 2021-08-18  
DE – PAUL HARTMANN AG  
Paul Hartmann-Strabe 12 · 89522 HEIDENHEIM, GERMANY  
AT – PAUL HARTMANN Ges.m.b.H. · 2355 Wiener Neudorf  
CH – IFF HARTMANN AG · 8212 Neuchâten

**Instructions for use****Product description**

PermaFoam Classic is a hydrophilic polymer with a three-dimensional foam structure which can effectively absorb wound exudate and maintain a moist wound environment. PermaFoam Classic Border also has an adhesive border coated with a polyacrylate adhesive.

**Composition**

- PU foam
- Polyacrylate
- PU film

**Indications**

PermaFoam Classic is a sterile, non-medicated single-use polyurethane foam dressing indicated for medium to highly exuding, acute and chronic wounds, such as venous ulcers (leg ulcers), pressure ulcers II-IV, diabetic foot ulcers, donor sites, abrasions, incisions.

**Contraindications**

PermaFoam Classic should not be used on patients with a known hypersensitivity to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Special instructions**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy:** see diagram 3  
**PermaFoam Classic Border:** see diagram 5

**Contraindications**

PermaFoam Classic is contraindicated on any patients who may be allergic to any of its ingredients.

**Precautions**

- Do not use the product in case of infected wounds.
- If redness or allergic reactions occur, please stop treatment with PermaFoam Classic.
- If the wound is caused by circulatory disorder or diabetic foot ulcer, please check the wound status on a daily basis.
- Please remove the product before applying radiation treatment, including X-ray, ultrasonic therapy, thermotherapy and microwave therapy.
- Do not use PermaFoam Classic in combination with hydrogen peroxide.

**Product disposal**

In order to minimize the risk of potential infection hazards, or environmental pollution, disposable components of PermaFoam Classic should follow disposal procedures according to applicable and local laws, regulations and infection prevention standards.

PermaFoam Classic can be used for up to 7 days. Please note that the frequency of dressing change depends on the condition of the wound and the level of exudate.

**PermaFoam Classic Sacral:** see diagram 1  
**PermaFoam Classic Concave:** see diagram 2  
**PermaFoam Classic Tracheostomy**

